arXiv:1507.02767v3 [math.DG] 2 Sep 2016

Unstable CMC spheres and outlying CMC spheres in AF 3-manifolds

Shiguang Ma*

September 5, 2016

Abstract

In this paper, we introduce a non-linear ODE method to construct CMC surfaces in Riemannian manifolds with symmetry. As an application we construct unstable CMC spheres and outlying CMC spheres in asymptotically Schwarzschild manifolds with metrics like $g_{ij} = (1 + \frac{1}{l})^2\delta_{ij} + O(l^{-2})$. The existence of unstable CMC spheres tells us that the stability condition in Qing-Tian's work [12] can not be removed generally.

1 Introduction

Constant mean curvature (CMC) surfaces are a kind of important submanifolds. In this paper we mainly focus on the CMC spheres in asymptotically flat (AF) 3-manifolds. First let's state some background works in this area.

In 1996, G. Huisken and S.T. Yau proved the existence of a foliation of stable CMC spheres in asymptotically Schwarzschid manifolds in [5]. They also proved, under certain radius condition, the uniqueness of the stable CMC spheres. In [12], J. Qing and G. Tian removed this radius condition, i.e. they proved that, outside certain compact subset, any stable CMC sphere which separates the compact part from infinity belongs to those constructed in [5]. There are many sequential works. Lan-hsuan Huang did similar work as Huisken and Yau in general AF manifolds with RT conditions (those are a series of asymptotically "odd" or "even" conditions). See [4]. In [9], Nerz considered the existence of CMC foliation in AF manifolds without RT conditions. The uniqueness results of Huang and Nerz also need radius conditions. In [6], I proved a uniqueness result under mild radius condition, which improved Huang's uniqueness result in [4] in some special case. In [7], I removed the radius condition in proving the uniqueness in AF manifolds with decay rate of the metic to be -1. Gang Tian and Andre Neves have also

---

*School of Mathematical Sciences and LPMC, Nankai University, Tianjin 300071, People's Republic of China, msgdyx8741@nankai.edu.cn.The author was supported by NSFC grant No.11301284, NSFG grant No. 11571185 and "Specialized Research Fund for the Doctoral Program of Higher Education", No.20120031120028.

considered similar existence and uniqueness problems in asymptotically hyperbolic manifolds in [10] and [11].

However, in all the above works, one has to assume stability of the CMC surfaces to prove the uniqueness (In Nerz's work he used an integral estimate on the mean curvature instead). It was asked by Professor Gang Tian that whether one can remove the stability condition in proving the uniqueness and if the answer is no generally, when we can do this. In [1] Simon Brendle prove that in a class of warped product metrics the only CMC spheres are umbilic. In particular, in Schwarzschild manifolds with positive mass, the only embedded CMC surfaces are spheres of symmetry. This theorem requires assumptions on neither the topology nor the stability of the surfaces. Then it is natural to consider asymptotically Schwarzschild manifolds with positive mass. In this paper without loss of generality we consider asymptotically Schwarzschild manifolds with only one asymptotically flat end. Suppose (M, ds^2) is a complete Riemannian manifold. For a compact subset $K \subset M$, we assume $M \backslash K$ is diffeomorphic to $\mathbb{R}^3 \backslash \bar{B}_1(0)$ and we assume $\{x_i\}_{i=1}^3$ are the standard coordinates of $\mathbb{R}^3$. Denote

$$l = \sqrt{x_1^2 + x_2^2 + x_3^3}.$$

Note that the Schwarzschild metric with mass 1 takes the form

$$g_{ij}^S = (1 + \frac{1}{2l})^4 \delta_{ij}$$

with δ_{ij} to be the standard Euclidean metric.

We use the cylindrical coordinates (x, r, θ) to calculate, namely,

$$\begin{cases} x_1 = x, \\ x_2 = r \cos\theta, \\ x_3 = r \sin\theta, \end{cases}$$

where $r = \sqrt{x_2^2 + x_3^2}$.

For $\lambda > 0$ and $p \in \mathbb{R}$, let $\phi(s) = \phi_{\lambda,p}(s), s \in [0, +\infty]$ be a smooth monotonic function such that

$$\phi(s) = \begin{cases} 1, & s \in [0, \lambda], \\ p, & s \in [2\lambda, +\infty], \end{cases} \tag{1}$$

and

$$\lambda|\phi'| + \lambda^2|\phi''| \leq C|p - 1|. \tag{2}$$

We consider asymptotically Schwarzschild metrics which take the following form on $\mathbb{R}^3 \backslash \bar{B}_1(0)$

$$ds^2(\partial_i, \partial_j) = \delta_{ij} + T_{ij}$$

where

$$\begin{cases} T_{rr} &= g_1(r, x) = \frac{2}{l} + \frac{1}{l^2}, \\ T_{\theta\theta} &= g_2(r, x) = r^2(\frac{2}{l} + \frac{1}{l^2}\phi_{\lambda,p}(|\frac{r}{x}|)), \\ T_{xx} &= g_3(r, x) = \frac{2}{l} + \frac{1}{l^2}, \\ T_{rx} &= T_{r\theta} = T_{x\theta} = 0. \end{cases} \tag{3}$$

Note that when $|\frac{r}{x}| = +\infty$, it should be understood as that $x = 0$.

Let

$$\chi(r,x) = \frac{1 + \frac{2}{l} + \frac{1}{l^2}\phi_{\lambda,p}(|\frac{r}{x}|)}{1 + \frac{2}{l} + \frac{1}{l^2}}.$$

So we have $r^2 + T_{\theta\theta} = r^2\chi(r,x)(1 + g_1(r,x))$.

It is obvious that for fixed λ, p, ds^2 is a smooth metric and

$$ds^2 = g^S + O(l^{-2}).$$

Remark. Note that

$$\begin{cases} x_2 &= r\cos\theta, \\ x_3 &= r\sin\theta. \end{cases}$$

Assume $x = x_1 > 0$. Let $ds^2 - (1 + \frac{1}{l})^2\delta_{ij} = S_{ij}$. Then by direct calculation we have

$$S_{ij,ij} - S_{ii,jj} = -\frac{\frac{2x\phi'(\frac{r}{x})}{r} + \phi''(\frac{r}{x})}{x^4}.$$

For any $0 < \alpha < \beta$.

$$\begin{aligned} &\int_{\alpha \le x_1 \le \beta} (S_{ij,ij} - S_{ii,jj}) dx_1 dx_2 dx_3 \\ =& -2\pi \int_\alpha^\beta \frac{1}{x^2} dx \int_\lambda^{2\lambda} (2\phi'(s) + s\phi''(s)) ds \\ =& 2\pi(\frac{1}{\beta} - \frac{1}{\alpha})(p - 1). \end{aligned}$$

So when $p > 1$, $S_{ij,ij} - S_{ii,jj}$ makes a negative contribution to the scalar curvature. Whether the scalar curvature of an asymptotically Schwarzschild manifold is positive is a very delicate thing. When the scalar curvature is negative, we have more flexibility while when the scalar curvature is positive, we have more rigidity. This can also be seen in [2] and [3].

Now we can state the main theorems of this paper.

Theorem 1.1. *Suppose $(M\backslash K, ds^2)$ is an asymptotically flat end with $ds^2(\partial_i, \partial_j) = \delta_{ij} + T_{ij}$ with T_{ij} given by (3). Then we can fix $\lambda > 0$ small and $p > 0$ large such that there is $\delta(\lambda, p) > 0$ such that for $0 < H < \delta(\lambda, p)$, there is an embedded CMC sphere Σ which is unstable, has mean curvature H and separates K from infinity. All the unstable CMC spheres constructed can be parameterized by its mean curvature H. If we denote the area of $\Sigma(H)$ by $|\Sigma(H)|$ and $l_0 = \inf_{x\in\Sigma} |x|$ then we have*

$$l_0 = O(H^{-1}),$$
$$|H^2|\Sigma(H)| - 48\pi| \le C(p)H.$$

This theorem answers Tian's question partially. In the case of [12] (hence in the case of [7]), the stability condition can not be removed.

In a more recent work [2] of S. Brendle and M. Eichmair, they constructed outlying CMC spheres which are stable in asymptotically Schwarzschild manifold with positive mass. By "outlying" they meant that the compact region bounded by Σ is disjoint from $B_{l_0(\Sigma)}(0)$. Our approach also works in such a setting.

Theorem 1.2. *There is an asymptotically Schwarzschild manifold* (M, g) *with* $m = 1$ *in which we can construct two sequences of embedded outlying CMC spheres which are parameterized by* Σ_n^1, Σ_n^2 *and a sequence of CMC spheres* Σ_n^3 *which separate the compact part from infinity, such that*

$$H(\Sigma_n^1) = H(\Sigma_n^2) = H(\Sigma_n^3)$$

and $\Sigma_n^1, \Sigma_n^3, \Sigma_n^2$ *share a common symmetry axis and* Σ_n^1, Σ_n^2 *are tangent to* Σ_n^3 *at the two poles of* Σ_n^3 *where* $x_2 = x_3 = 0$.

This result is a little weaker than that of [2] in that we do not discuss the stability of the surfaces.

The main technique of this paper is direct analysis of a nonlinear ODE which comes from the study of Delaunay type CMC surfaces. In a recent work [8], Frank Pacard and I proved that along any non-degenerate closed embedded geodesic in a 3-manifold, one can construct constant mean curvature surfaces of Delaunay type of arbitrarily small size. It was asked by Frank Pacard that if we can deform the neck size of a Delaunay type CMC surface along a closed geodesic to develop a singularity (which can be regarded as the boundary of the moduli of CMC surfaces). If one can do this and prove that the singularity is removable in certain sense, then one can construct CMC clusters in Riemannian manifolds. In certain sense, this paper is also related to this problem. The unstable CMC spheres constructed are in fact CMC clusters of "three bubbles" in asymptotically Schwarzschild manifold. The author believes that this ODE method can be used to construct many examples. Also we can use it to treat the question raised by Pacard in the case that the metric has rotational symmetry along the geodesic.

The main contributions can be summarized as follows: The meridian curve of the CMC surfaces of revolution is determined by an ODE of the following type

$$\begin{cases} g_a''(y) & -\frac{1}{g_a(y)}(1 + g_a'(y)^2) + (2 + \rho)(1 + g_a'(y)^2)^{\frac{3}{2}} = 0, \\ g_a(0) & = a \in [0.95, 1.05], \\ g_a'(0) & = 0. \end{cases}$$

where the expression of $\rho = \rho(H, g_a, g_a', y)$ is given by (7) and $|\rho| \leq CH$ holds, where H is the mean curvature, which is assumed to be a small positive number. If we neglect ρ, when $a < 1$, the solution is the meridian curve of Delaunay unduloid in $\mathbb{R}^3$. As $a \to 1^-$ the solution develops singularities at each integer at the same time. And the Delaunay surface converges to infinitely many spheres each one of which meets its two neighbors at its two poles. If we do not neglect ρ, for a fixed $a < 1$, when H is very small, the solution looks similar to $\rho = 0$

case. Look at the graph below. The difference is that the solution is no longer exactly periodic. When a goes up, the solution $g_a(y), y \in [0,5]$ will develop a singularity. The singularity will develop at z_1 or z_2. To study this, we use the Delaunay parameter function

$$\tau(g_a(y), g_a'(y)) = -g_a^2(y) + \frac{g_a(y)}{\sqrt{1+g_a'(y)^2}}.$$



Before a singularity develops, $\tau(g_a(z_1), g_a'(z_1)), \tau(g_a(z_2), g_a'(z_2)) > 0$. When a singularity develops, $g_a \to 0, g_a' \to \infty$, so $\tau \to 0$. So we will compare $\tau(g_a(z_1), g_a'(z_1))$ with $\tau(g_a(z_2), g_a'(z_2))$. Note that we have

$$\frac{d}{dy}\tau(g_a(y), g_a'(y)) = g_a g_a' \rho.$$

So

$$\tau(g_a(z_2), g_a'(z_2)) - \tau(g_a(z_1), g_a'(z_1)) = \int_{z_1}^{z_2} g_a g_a' \rho dy.$$

So we have to study the function ρ. In our case, after some more calculations and arguments, we know (7) is reduced to

$$\rho = \frac{H}{d} + \frac{H(g_a - yg_a')}{d^3\sqrt{1+g_a'^2}} + \frac{H^2(\phi-3)(g_a - yg_a')}{4d^3\sqrt{1+g_a'^2}} - \frac{H^2\phi'(y+g_a g_a')}{8d^2y^2\sqrt{1+g_a'^2}} + C(p,\lambda)O(H^3)$$

where ϕ is given by (1). Note that ϕ has two parameters λ, p. At first glance, ϕ only appears in H^2 terms, which is not dominant. Here our key observation is that when H is very small

$$\int_{z_1}^{z_2} g_a g_a' (\frac{H}{d} + \frac{H(g_a - yg_a')}{d^3\sqrt{1+g_a'^2}})dy = O(H^2)$$

which is verified in Lemma 4.7. So when $|p|$ is very large, the third term becomes dominant. So we have freedom to choose the place where the singularity develops, by choosing proper p and λ. Despite the singularity, the surface of revolution is still smooth from Lemma 4.9. If the singularity develops at z_2, we can get the "three-bubble" like CMC sphere by applying symmetric extension. Still, we can choose λ and p properly such that the singularity develops simultaneously at z_1 and z_2. Then we can construct outlying CMC spheres.

The paper is organized as follows. In Section 2, we derive the ODE which is satisfied by the meridian curve of a surface of revolution with constant mean curvature. In Section 3, we analyze this ODE and prove the existence of the solution. In Section 4, we deform the Delaunay parameter until the singularity appears. And we can analyze the behavior of the solution at the singular point. In Section 5, we prove the two theorems.

2 The mean curvature

Suppose a surface of revolution $\Sigma \subset M$ is defined by

$$r = f(x)$$

where $f(x)$ is a smooth positive function. We are going to calculate the mean curvature of such a surface.

We assume (s, θ) to be the coordinate on the surface which satisfies

$$\frac{\partial}{\partial s} = \frac{\partial}{\partial x} + \frac{\partial f}{\partial x}\frac{\partial}{\partial r}, \frac{\partial}{\partial \theta} = x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2}.$$

We have

$$\begin{cases} g(\partial_s, \partial_s) & = 1 + f'(x)^2 + g_3 + f'(x)^2 g_1, \\ g(\partial_\theta, \partial_\theta) & = r^2 + g_2. \end{cases}$$

The unit normal vector on Σ is

$$V = V_r \partial_r + V_x \partial_x$$

where

$$\begin{cases} V_r & = \frac{1}{\sqrt{1+g_1}\sqrt{1+\frac{1+g_1}{1+g_3}f'^2}}, \\ V_x & = -f' \frac{\sqrt{1+g_1}}{(1+g_3)\sqrt{1+\frac{1+g_1}{1+g_3}f'^2}}. \end{cases}$$

We use V_e to represent the normal vector in Euclidean metric, i.e.

$$V_e = \frac{\partial_r - f'\partial_x}{\sqrt{1+f'^2}}.$$

We use $< \cdot, \cdot >$ to represent the inner product in the metric g and "$\cdot$" to represent the inner product in the Euclidean metric. The mean curvature

$$H(\Sigma) = g^{ss} < \nabla_{\partial_s} V, \partial_s > + g^{\theta\theta} < \nabla_{\partial_\theta} V, \partial_\theta >$$

$$\begin{aligned} < \nabla_{\partial_s} V, \partial_s > = & < (\partial_s V_r)\partial_r, \partial_s > + < V_r \nabla_{\partial_s} \partial_r, \partial_s > \\ & + < (\partial_s V_x)\partial_x, \partial_s > + < V_x \nabla_{\partial_s} \partial_x, \partial_s > . \end{aligned}$$

By direct calculations we have

$$\begin{aligned}
<(\partial_s V_r)\partial_r, \partial_s> &= (\partial_s V_r) f'(1+g_1),\\
<(\partial_s V_x)\partial_x, \partial_s> &= (\partial_s V_x)(1+g_3),\\
<V_r \nabla_{\partial_s}\partial_r, \partial_s> &= \frac{V_r}{2}(\partial_r g_{xx} + f'^2 \partial_r g_{rr}),\\
<V_x \nabla_{\partial_s}\partial_x, \partial_s> &= \frac{V_x}{2}(\partial_x g_{xx} + f'^2 \partial_x g_{rr}),
\end{aligned}$$

$$\begin{aligned}
<\nabla_{\partial_\theta} V, \partial_\theta> &= V_r <\nabla_{\partial_\theta}\partial_r, \partial_\theta> + V_x <\nabla_{\partial_\theta}\partial_x, \partial_\theta>\\
&= \frac{1}{2}(V_r \partial_r g_{\theta\theta} + V_x \partial_x g_{\theta\theta}).
\end{aligned}$$

So we get

$$\begin{aligned}
H(\Sigma) = {} & (1+f'(x)^2 + g_3 + f'(x)^2 g_1)^{-1}[(\partial_s V_r) f'(1+g_1) + (\partial_s V_x)(1+g_3)\\
& + \frac{V_r}{2}(\partial_r g_{xx} + f'^2 \partial_r g_{rr}) + \frac{V_x}{2}(\partial_x g_{xx} + f'^2 \partial_x g_{rr})]\\
& + (r^2 + g_2)^{-1} \frac{1}{2}(V_r \partial_r g_{\theta\theta} + V_x \partial_x g_{\theta\theta}).
\end{aligned}$$

From (3), the expression of $H(\Sigma)$ can be reduced to

$$\begin{aligned}
H(\Sigma) = & \frac{1}{2}(1+g_1)^{-\frac{3}{2}}(1+f'^2)^{-\frac{3}{2}}(-2(1+g_1)f''\\
& + (1+f'^2)(\partial_r g_1 - f'\partial_x g_1)\\
& + (r^2+g_2)^{-1}(1+g_1)(1+f'^2)(2r + \partial_r g_2 - f'\partial_x g_2))\\
= & -(1+g_1)^{-\frac{1}{2}}(1+f'^2)^{-\frac{3}{2}} f'' - 2(V_e \cdot x')(l^{-3} + l^{-4})(1+g_1)^{-\frac{3}{2}}\\
& + (1+g_1)^{-\frac{1}{2}}(1+f'^2)^{-\frac{1}{2}} f^{-1} + \frac{1}{2}(1+g_1)^{-\frac{1}{2}} \chi(r,x)^{-1} V_e(\chi).
\end{aligned}$$

where $x' = (x, f(x)\cos\theta, f(x)\sin\theta)$ and $V_e(\chi)$ is the derivative of $\chi(r,x)$ in the direction $V_e = \frac{\partial_r - f'\partial_x}{\sqrt{1+f'^2}}$.

Now we have

$$\frac{d^2 f}{dx^2} - \frac{1}{f}(1 + (\frac{df}{dx})^2) + (H + \tilde{\rho})(1 + (\frac{df}{dx})^2)^{\frac{3}{2}} = 0, \tag{4}$$

where

$$\tilde{\rho} = \frac{H}{l} + 2(V_e \cdot x')(l^{-3} + l^{-4})(1+g_1)^{-1} - \frac{1}{2}\chi^{-1} V_e(\chi).$$

Since we want to get constant mean curvature surfaces, we assume H to be a small positive constant. Denote

$$g(y) = \frac{H}{2} f(\frac{2}{H} y) = \frac{H}{2} f(x).$$

We have

$$g'(y) = f_x(\frac{2}{H}y), g''(y) = \frac{2}{H}f_{xx}(\frac{2}{H}y),$$

So we have

$$g''(y) - \frac{1}{g(y)}(1 + g'(y)^2) + (2 + \rho)(1 + g'(y)^2)^{\frac{3}{2}} = 0 \tag{5}$$

where

$$\rho = \frac{2}{H}\tilde{\rho}.$$

We want to study ODE (5) instead of (4).

3 The analysis of the ODE

Consider

$$\begin{cases} g''(y) - \frac{1}{g(y)}(1 + g'(y)^2) + (2 + \rho)(1 + g'(y)^2)^{\frac{3}{2}} = 0, \\ g(0) = a, \\ g'(0) = 0. \end{cases} \tag{6}$$

We assume $a \in [0.9, 1.1]$ and denote the solution by $g_a(y)$. Let us assume that $H > 0$ is very small and the real number p is fixed. Let $d = \frac{H}{2}l$. Then we have

$$\begin{aligned} \rho =& \frac{H}{d} + (\frac{H}{2} + \frac{H^2}{4d})\frac{g_a - yg_a'}{d^3(1 + g_1)\sqrt{1 + g_a'^2}} \\ &+ (\frac{H}{2} + \frac{H^2\phi}{4d})\frac{g_a - yg_a'}{d^3(1 + \frac{H}{d} + \frac{\phi H^2}{4d^2})\sqrt{1 + g_a'^2}} \\ &- \frac{H^2}{8d^2}\frac{\phi'}{(1 + \frac{H}{d} + \frac{\phi H^2}{4d^2})}\frac{y + g_a g_a'}{y^2\sqrt{1 + g_a'^2}}. \end{aligned} \tag{7}$$

In the following analysis, by a constant C (or $C_i, i = 1, 2, \cdots$) we mean a general (or a particular) uniform positive constant which does not depend on H, a,p, λ or y. We use symbols $C(H), C(p)$ to denote constants which depend on H, p, etc.

First we have the local existence. Let A_1, A_2, A_3, A_4 be four positive constant. Let's denote a domain in the phase space

$$\{(g, g'); A_2 \leq g \leq A_1, -A_4 \leq g' \leq A_3\}$$

as $D(A_1, A_2, A_3, A_4)$.

Lemma 3.1. *Suppose $g_a(y), y \in [\alpha, \beta], 0 \leq \alpha \leq \beta$ solves (6). And for some $A_i, i = 1, \cdots, 4$, $(g_a, g_a') \in D(A_1, A_2, A_3, A_4), y \in [\alpha, \beta]$. Then there is $\delta > 0$ such that the solution g_a can be extended to $[\alpha, \beta + \delta)$.*

Proof. Consider a system that is equivalent to (6)

$$\begin{cases} g_a' &= w_a, \\ w_a' &= \frac{1}{g_a}(1+w_a^2) - (2+\rho)(1+w_a^2)^{\frac{3}{2}}. \end{cases}$$

When $(g_a, w_a) \in D(A_1, A_2, A_3, A_4)$ and $y \in [\alpha, \beta]$, we know d has lower bound A_2. The right hand side has uniform bound and uniform Lipschitz constant with respect to g_a, w_a. From the local existence of ODE system, we can prove this lemma.

□

Then we have a “closed” property for the solution.

Lemma 3.2. *Suppose* $g_a(y), y \in [\alpha, \beta), 0 \le \alpha < \beta < +\infty$ *solves (6). And for some* $A_i > 0, i = 1, \cdots, 4$, $(g_a, g_a') \in D(A_1, A_2, A_3, A_4), y \in [\alpha, \beta)$. *If*

$$\lim_{y \to \beta^-} g_a(y), \lim_{y \to \beta^-} g_a'(y)$$

exist and are finite, then $g_a(y)$ *(as a solution to (6)) can be extended to* $y = \beta$. *In particular, if* $g_a(y)$ *and* $g_a'(y)$ *are both monotonic and bounded,* g_a *can be extended to* $y = \beta$.

Proof. If for some $A_i, i = 1, \cdots, 4$, $(g_a, g_a') \in D(A_1, A_2, A_3, A_4), y \in [\alpha, \beta)$ and

$$\lim_{y \to \beta^-} g_a(y), \lim_{y \to \beta^-} g_a'(y)$$

exist, then from (6) and (7) we have

$$\lim_{y \to \beta^-} g_a''(y)$$

exists. By defining the values of g_a, g_a', g_a'' on β in the obvious way, we can prove this lemma.

□

Lemma 3.3. *Suppose* $g_a(y) > 0, y \in I$ *solves (6) where* I *is an interval* $[\alpha, \beta]$ *or* $[\alpha, \beta)$, $0 \le \alpha < \beta$. *In the case of* $I = [\alpha, \beta)$, $g_a(\beta)$ *and* $g_a'(\beta)$ *should be understood as a limit which is assumed to exist.*

Then

1. *If* $g_a(y)$ *is monotonic in* y *and* $0 < C^{-1} < |g_a'| < C$ *when* $y \in I$, *then*

$$C^{-1}|g_a(\beta) - g_a(\alpha)| \le |\beta - \alpha| \le C|g_a(\beta) - g_a(\alpha)|.$$

2. *If* $g_a'(y)$ *is monotonic in* y *and* $k_a = g_a\sqrt{1 + g_a'(y)^2}$ *satisfies that* $k_a - \frac{1}{2}$ *does not change sign and*

$$C^{-1} \le |k_a - \frac{1}{2}| \le C, y \in I.$$

And we assume that $|\rho| \leq CH$. Then when H is small we have

$$C^{-1}\inf_{y\in I} g_a \leq \frac{|\beta-\alpha|}{|\arctan g_a'(\beta) - \arctan g_a'(\alpha)|} \leq C\sup_{y\in I} g_a.$$

Proof. We need only prove the case $I = [\alpha, \beta]$. The case $I = [\alpha, \beta)$ follows by considering $[\alpha, \beta_i]$ with $\beta_i \to \beta$.

The first one follows form

$$|\beta - \alpha| = |\int_{g_a(\alpha)}^{g_a(\beta)} \frac{dg_a}{g_a'}|$$

and the second one follows from the fact that ρ is small and

$$|\beta - \alpha| = |\int_{g_a'(\alpha)}^{g_a'(\beta)} \frac{dg_a'}{g_a''}| = |\int_{g_a'(\alpha)}^{g_a'(\beta)} \frac{g_a dg_a'}{(1+g_a'^2)(1-(2+\rho)k_a)}|.$$

□

Now we will prove that for the solution of (6), $d = \sqrt{y^2+g_a^2}$ has uniform positive lower bound independent of H and $a \in [0,9,1,1]$. This is an important estimate. Only with this estimate can we know that $|\rho| \leq CH$.

Lemma 3.4. *We can choose $\lambda > 0$ small and $\delta(p,\lambda) > 0$ such that when $H < \delta(p,\lambda)$, for all $a \in [0.9, 1.1]$, as long as the solution of (6) can be extended when $y > 0$,*

$$\begin{aligned} d = \sqrt{g_a^2+y^2} &\geq C > 0, \\ |\frac{1}{8d^2}\frac{\phi'}{(1+\frac{H}{d}+\frac{\phi H^2}{4d^2})}\frac{y+gg'}{y^2\sqrt{1+g_a'^2}}| &\leq \frac{C|p-1|}{\lambda}, \\ |\rho| &\leq \hat{C}H. \end{aligned} \tag{8}$$

Proof. First let us pretend that $\phi = p$ holds everywhere. So $\phi' = 0$ and

$$\begin{aligned} \rho =& \frac{H}{d} + \frac{H}{2}\frac{g_a - yg_a'}{d^3(1+g_1)\sqrt{1+g_a'^2}} + \frac{H^2}{4}\frac{g_a-yg_a'}{d^4(1+g_1)\sqrt{1+g_a'^2}} \\ &+ \frac{H}{2}\frac{g_a-yg_a'}{d^3(1+\frac{H}{d}+\frac{\phi H^2}{4d^2})\sqrt{1+g_a'^2}} + \frac{H^2}{4}\frac{p(g_a-yg_a')}{d^4(1+\frac{H}{d}+\frac{\phi H^2}{4d^2})\sqrt{1+g_a'^2}}. \end{aligned}$$

Let $D_a(y)$ solves

$$\begin{cases} D''(y) - \frac{1}{D(y)}(1+D'(y)^2) + 2(1+D'(y)^2)^{\frac{3}{2}} = 0, \\ D(0) = a, \\ D'(0) = 0, \end{cases} \tag{9}$$

where $a \in [0.8, 1.2]$.

We state some facts of $D_a(y)$ without proof. There are $\delta_0, C_1, C_2, C_3 > 0$ such that,

1. For each $a \in [0.8, 1.2]$, the solution of (9) exists on $[0, \delta_0]$.
2. $R_1 = \{(D_a(y), \frac{d}{dy}D_a(y)); a \in [0.8, 1.2], y \in [0, \delta_0]\}$ is trapped in the region

$$R_2 = \{(\xi, \eta); C_1 \leq \xi \leq 1.2, -C_2 \leq \eta \leq 0\}.$$

3. When $a \in [0.8, 1.2]$, the solution $D(y), y \in [0, \delta_0]$ satisfies $D\sqrt{1 + D'^2} \geq \frac{2}{3}$.
4. For each $a \in [0.8, 1.2]$, $D_a(0) - D_a(\delta_0) > C_3$ and $D'_a(\delta_0) < -C_3$.

We may draw a graph in the phase space to illustrate this. In the graph below, $\delta_0 = 0.75$. We can choose $C_1 = 0.5, C_2 = 2.4$. $D_a(0) - D_a(\delta_0)$ and $-D'_a(\delta_0)$ are positive continuous functions of $a \in [0, 8, 1.2]$. So C_3 can be chosen properly small and positive.



Let R_1 be the closure of a bounded domain whose boundary contains 4 components, $\Gamma_1, \Gamma_2, \Gamma_3, \Gamma_4$.

$$\Gamma_1 = \{(D_a(y), \frac{d}{dy}D_a(y)); a = 1.2, y \in [0, \delta_0]\},$$

$$\Gamma_2 = \{(D_a(y), \frac{d}{dy}D_a(y)); a \in [0.8, 1.2], y = \delta_0\},$$

$$\Gamma_3 = \{(D_a(y), \frac{d}{dy}D_a(y)); a = 0.8, y \in [0, \delta_0]\},$$

$$\Gamma_4 = \{(D_a(y), \frac{d}{dy}D_a(y)); a \in [0.8, 1.2], y = 0\}.$$

Now we consider the orbit (g_a, g'_a) of (6). For each $a \in [0, 9, 1.1]$, $(g_a(0), g'_a(0)) \in \Gamma_4$. From Lemma 3.1, the solution of (6) can be extended a little to $y > 0$. Then the orbit will be extended into the interior of R_1. As long as the orbit keeps in R_1, we know $d \geq C_1$. So there are $\delta(p) > 0, C > 0$ such that when $0 < H < \delta(p)$,

$$|\rho| \leq CH. \tag{10}$$

If $\{(g_a(y), g_a'(y)); y \in [0, \delta_0]\} \subset R_1$, we have $d = \sqrt{y^2 + d^2} \geq \min\{\delta_0, C_1\}$ as long as the solution exists. If it is not the case then we assume $y' \in (0, \delta_0)$ is the first time (g_a, g_a') touches ∂R_1. We claim that for H sufficiently small, $(g_a(y'), g_a'(y')) \in \Gamma_2$. For this we consider the Delaunay parameter function

$$\tau(g_a, g_a') = -g_a^2 + \frac{g_a}{\sqrt{1 + g_a'^2}}.$$

One can regard it as the first integral of (9). When $y \in [0, y']$

$$\frac{d}{dy}\tau = g_a g_a' \rho. \tag{11}$$

Note that $|g_a g_a'| \leq 1.2C_2$. From (10), by choosing H even smaller, we have for each $a \in [0, 9, 1.1]$, as long as $(g_a, g_a') \in R_1$, (g_a, g_a') keeps in a small neighborhood of (D_a, D_a'). So (g_a, g_a') has no chance to touch Γ_1 or Γ_3. From

$$g_a\sqrt{1 + g_a'^2} \geq \frac{2}{3} \tag{12}$$

and the fact that ρ is small, we can deduce

$$g_a''(y) = \frac{1}{g_a(y)}(1 + g_a'(y)^2)(1 - (2 + \rho)g_a(y)(1 + g_a'(y)^2)^{\frac{1}{2}}) \tag{13}$$

is always negative, which implies both g_a and g_a' are monotonically decreasing when $y \in [0, y']$. From Lemma 3.3, we have

$$C(C_1, C_2, C_3)^{-1} \leq y' \leq C(C_1, C_2, C_3).$$

So if we choose $2\lambda \leq C(C_1, C_2, C_3)^{-1}C_1$ and define ϕ as (1), it will have no influence on the analysis above. And we know

$$d \geq \min\{C(C_1, C_2, C_3)^{-1}, C_1, \delta_0\}.$$

Note that when $\phi' \neq 0$, there hold $y \geq C(C_1, C_2, C_3)^{-1}$ and $|\frac{g}{y}| \leq 2\lambda$. So from (2)

$$\begin{aligned} &|\frac{1}{8d^2}\frac{\phi'}{(1 + \frac{H}{d} + \frac{\phi H^2}{4d^2})}(1 + g'^2)^{-\frac{1}{2}}\frac{y + gg'}{y^2}| \\ \leq &\frac{C|p - 1|}{\lambda}. \end{aligned}$$

So from

$$\frac{|g_a - yg_a'|}{\sqrt{1 + g_a'^2}\sqrt{y^2 + g_a^2}} \leq 1$$

we can choose $\delta(p, \lambda)$ small such that $|\rho| \leq \hat{C}H$.

$\square$

Now we can expand ρ in terms of H. For $0 < H < \delta(p,\lambda)$,

$$\rho = \frac{H}{d} + \frac{H(g_a - yg_a')}{d^3\sqrt{1+g_a'^2}} + \frac{H^2(\phi-3)(g_a - yg_a')}{4d^3\sqrt{1+g_a'^2}} - \frac{H^2\phi'(y+g_ag_a')}{8d^2y^2\sqrt{1+g_a'^2}} + C(p,\lambda)O(H^3).$$

From now on we will always assume that $\delta(p,\lambda)$ is so small that $|\rho| \le \hat{C}H$.

It is obviously that exactly one of the following three will happen to $g_a(y)$,

(H1): The solution $g_a(y)$ can be extended at least two times such that $g_a(y) \in (0,\frac{1}{2}), g_a'(y) = 0$;

(H2): The solution $g_a(y)$ can be extended only one time such that $g_a(y) \in (0,\frac{1}{2}), g_a'(y) = 0$;

(H3): The solution $g_a(y)$ cannot be extended to a point y such that $g_a(y) \in (0,\frac{1}{2}), g_a'(y) = 0$.

Let's draw some pictures of the phase space to illustrate the three cases.



H1 happens



H2 happens



H3 happens

Lemma 3.5. *There is a uniform $C_4 > 0$ such that if $g_a(y)$ can be extended to some $y_1 > 0$ such that $0 < g_a(y_1) < C_4, g_a'(y_1) < 0$ and $g_a''(y_1) = 0$, then there is $y_2 > y_1$ such that $g_a(y_2) \in (0, \frac{1}{2})$ and $g_a'(y_2) = 0$. Moreover, in $[y_1, y_2]$, $g_a(y)$ is monotonically decreasing and $g_a'(y)$ is monotonically increasing. (This can be seen in the case "H1" and "H2".)*

Proof. From Lemma 3.1, we know that the solution can be extended beyond y_1 for a small interval. We want to show that, when $y > y_1$, as long as the solution exists and $g_a > 0$ and $g_a' < 0$, we have $g_a'' > 0$. As before we denote

$$k_a(y) = g_a\sqrt{1 + g_a'^2}.$$

Note that $|\rho| \leq \hat{C}H$. If $k_a(y) < \frac{1}{2+\hat{C}H}$, we have

$$g_a''(y) = \frac{1}{g_a}(1 + g_a'^2)(1 - (2 + \rho)k_a) > 0.$$

From easy calculation

$$\frac{dk_a(y)}{dy} = g_a'(1 + g_a'^2)^{\frac{1}{2}} + g_a(1 + g_a'^2)^{-\frac{1}{2}}g_a'g_a''. \tag{14}$$

When $g_a > 0, g_a' < 0$, we have

$$\begin{aligned}\frac{dk_a}{dg_a} &= \frac{dk_a(y)}{dy}\frac{1}{g_a'} = (1 + g_a'^2)^{\frac{1}{2}} + g_a(1 + g_a'^2)^{-\frac{1}{2}}g_a'' \\ &= \frac{k_a(2 - (2 + \rho)k_a)}{g_a}.\end{aligned}$$

So we have

$$\frac{dk_a}{k_a(2 - (2 + \rho)k_a)} = \frac{dg_a}{g_a}. \tag{15}$$

When $y = y_1$, from $g_a''(y_1) = 0$, we have $2 - (2 + \rho(y_1))k_a(y_1) = 1$. So as long as the solution can be extended when $y > y_1$ and $g_a > 0, g_a' < 0$ we have k_a is monotonically decreasing. So once $k_a(y) < \frac{1}{2+\hat{C}H}$, $g_a''(y) > 0$ holds as long as $y > y_1, g_a > 0, g_a' < 0$.

The problem is whether we still have $g_a''(y) > 0$, when $y > y_1$ and $\frac{1}{2+\rho(y_1)} \geq k_a \geq \frac{1}{2+\hat{C}H}$. We choose $0 < C_4 < \frac{1}{\sqrt{2}(2+\hat{C}H)}$. When $0 < g_a(y) \leq g_a(y_1) \leq C_4$, $k_a = g_a(y)\sqrt{1 + g_a'^2(y)} \geq \frac{1}{2+\hat{C}H}$ and $g_a'(y) < 0$ we have

$$\sqrt{1 + g_a'^2(y)} \geq \sqrt{2}$$

which implies $g_a'(y) \leq -1$.

From direct calculations, $\exists$ a uniform constant $C > 0$, such that when H is sufficiently small

$$\begin{aligned}|\rho'| &\leq (\frac{H}{d^3}y + \frac{H}{d^3}g_a|g_a'|)(1 + \frac{|g_a''|}{(1 + (g_a')^2)^{\frac{3}{2}}}) + C(p, \lambda)H^2(1 + |g_a'|) \\ &\leq CH(1 + |g_a'|).\end{aligned} \tag{16}$$

From

$$\frac{d}{dy}((2+\rho)k_a) = \rho' k_a + (2+\rho)k_a'$$

and (14)(16) we have

$$\frac{d}{dg_a}((2+\rho)k_a) \geq \frac{1}{2}\frac{k_a}{g_a}(2+\rho - |\frac{2CH(1+g_a')g_a}{g_a'}|). \tag{17}$$

When $y > y_1, k_a \geq \frac{1}{2+\hat{C}H}$, from

$$g_a' \leq -1$$

we have, for H sufficiently small, $\frac{d}{dg_a}((2+\rho)k_a)$ is positive. (Here we need the estimate $g_a' \leq -1$. We use C_4 for this technical reason.) So when $y > y_1$ and $\frac{1}{2+\rho(y_1)} \geq k_a \geq \frac{1}{2+\hat{C}H}$, we have $(2+\rho)k_a$ monotonically decreasing as y increases. So when $y > y_1$,

$$g_a''(y) = \frac{1}{g_a}(1+g_a'^2)(1-(2+\rho)k_a) > 0.$$

By now we have known, when $y > y_1$, as long as the solution exists and $0 < g_a \leq C_4$ and $g_a' < 0$, g_a is monotonically decreasing and g_a' is monotonically increasing. Now we prove that there is a positive lower bound for g_a. We assume the solution can be extended from y_1 to $\tilde{y}_2$ with $g_a(\tilde{y}_2) > 0$ and $g_a'(\tilde{y}_2) < 0$. From (15) we know when $y \in [y_1, \tilde{y}_2]$

$$\int_{g_a(y)}^{g_a(y_1)} \frac{dg_a}{g_a} = \int_{k_a(y)}^{k_a(y_1)} \frac{dk_a}{k_a(2-(2+\rho)k_a)} \leq \int_{k_a(y)}^{k_a(y_1)} \frac{dk_a}{k_a(2-(2+\hat{C}H)k_a)}.$$

So we have

$$\ln\frac{g_a(y_1)}{g_a(y)} \leq (\frac{1}{2}\ln\frac{(2+\hat{C}H)k_a}{2-(2+\hat{C}H)k_a})|_{k_a(y)}^{k_a(y_1)},$$

which implies

$$k_a(y) \leq C(\frac{g_a(y)}{g_a(y_1)})^2.$$

From

$$\frac{dg_a'}{dg_a} = \frac{g_a''}{g_a'} = \frac{(1+(g_a')^2)[1-(2+\rho)k_a]}{g_a g_a'}$$

we have

$$\begin{aligned}\int_{g_a'(\tilde{y}_2)}^{g_a'(y_1)} \frac{g_a' dg_a'}{1+(g_a')^2} &= \int_{g_a(\tilde{y}_2)}^{g_a(y_1)} \frac{1-(2+\rho)k_a}{g_a} dg_a \\ &\geq \int_{g_a(\tilde{y}_2)}^{g_a(y_1)} \frac{1-C(\frac{g_a(y)}{g_a(y_1)})^2}{g_a(y)} dg_a(y)\end{aligned}$$

which implies

$$\frac{1}{2}\ln(1+g_a'^2)|_{g_a'(\tilde{y}_2)}^{g_a'(y_1)} \geq \ln\frac{g_a(y_1)}{g_a(\tilde{y}_2)} - \frac{C}{2}g_a(y_1)^{-2}(g_a(y_1)^2 - g_a(\tilde{y}_2)^2).$$

So we have

$$g_a(\tilde{y}_2) \geq \mathrm{e}^{-C}(1+g_a'(y_1)^2)^{-\frac{1}{2}}g_a(y_1).$$

Suppose y_2 is the supremum of the values $\tilde{y}_2$ until which the solution can be extended and $g_a(\tilde{y}_2) > 0, g_a'(\tilde{y}_2) < 0$. From Lemma 3.3, we can prove that $\tilde{y}_2$ is bounded (We can divide $g_a, y \in [y_1, \tilde{y}_2]$ into two parts. On the first part g_a is bounded, $g_a' < -\frac{1}{2}$ and on the second part $-\frac{1}{2} < g_a' < 0$ and k_a is away from $\frac{1}{2}$), hence y_2 is finite.

From Lemma 3.2 and the monotonicity of g_a and g_a' we know g_a can be extended to y_2 and

$$\begin{aligned} &g_a(y_2) \geq \mathrm{e}^{-C}(1+g_a'(y_1)^2)^{-\frac{1}{2}}g_a(y_1),\\ &g_a'(y_2) \leq 0. \end{aligned}$$

So we must have $g_a'(y_2) = 0$, or there would be a contradiction with the fact that y_2 is the supremum of $\tilde{y}_2$. Because $C_4 < \frac{1}{2}$, we have $0 < g_a(y_2) < \frac{1}{2}$.

□

Lemma 3.6. *When $a \in [0.9, 1.1]$, if (H3) happens, then $\exists y_1' \in (0, +\infty)$ such that $g_a(y)$ can be extended until $(0, y_1')$ but not until $(0, y_1']$. Moreover, in $(0, y_1')$, $g_a(y)$ and $g_a'(y)$ are monotonically decreasing and*

$$\lim_{y\to y_1'^-} g_a(y) \geq 0, \lim_{y\to y_1'^-} g_a'(y) = -\infty.$$

Proof. When $a \in [0.9, 1.1]$, we claim, if (H3) happens, when $y > 0$, as long as the solution can be extended, $g_a(y) > 0, g_a'(y) < 0, g_a''(y) < 0$.

First we prove this claim. When $y \in (0, \delta)$ for a small δ, $g_a(y) > 0, g_a'(y) < 0, g_a''(y) < 0$ hold. If the claim were false then there would be some $y_1 > 0$ such that

$$g_a(y_1) > 0, g_a'(y_1) < 0, g_a''(y_1) = 0$$

and in $(0, y_1)$, $g_a(y) > 0, g_a'(y) < 0, g_a''(y) < 0$ hold. From the monotonicity of g_a we have

$$\frac{d\tau}{dg_a} = \frac{d\tau}{dy}\frac{1}{g_a'} = g_a\rho. \tag{18}$$

Since g_a is bounded, there is $C > 0$ such that

$$|\tau(g_a(y_1), g_a'(y_1)) - (a - a^2)| = |\tau(g_a(y_1), g_a'(y_1)) - \tau(a, 0)| \leq CH \tag{19}$$

$$g_a(y_1) = \sqrt{\frac{\tau(g_a(y_1), g_a'(y_1))}{\frac{1}{k_a(y_1)} - 1}} = \sqrt{\frac{a - a^2 \pm CH}{\frac{1}{k_a(y_1)} - 1}}.$$

Note that $k_a(y_1) \in [\frac{1}{2+\hat{C}H}, \frac{1}{2-\hat{C}H}]$. So we can choose H sufficiently small such that $g_a(y_1) \leq \sqrt{\frac{0.9-0.9^2+CH}{1-CH}} \approx 0.3$. One can choose C_4 in Lemma 3.5 such that $\sqrt{\frac{0.9-0.9^2+CH}{1-CH}} \leq C_4 \leq \frac{1}{2(\sqrt{2}+CH)}$. So $g_a(y_1) \leq C_4$. From Lemma 3.5 one can find a contradiction with (H3). So the claim is true.

Now we prove the lemma. Suppose the solution can be extended to $[0, y^*]$. From $g_a(y^*) > 0, g_a'(y^*) < 0, g_a''(y^*) < 0$ and Lemma 3.1, the solution can be extended beyond y^*. So we know the maximal interval in which the solution exists should look like $[0, y_1')$ with y_1' possibly be $+\infty$. To see that y_1' is finite, we just prove that when $-\frac{1}{2} < g_a' < 0$, k_a is away from $\frac{1}{2}$. Then we can apply Lemma 3.3, by dividing the orbit into two parts. Note that

$$k_a = g\sqrt{1+g'^2} = \frac{1 \pm \sqrt{1-4\tau(1+g'^2)}}{2}.$$

So we have

$$|k_a - \frac{1}{2}| = \frac{\sqrt{1-4\tau(1+g'^2)}}{2}.$$

From (19) and $g'^2 \leq \frac{1}{4}$ we have

$$|k_a - \frac{1}{2}| \geq \frac{\sqrt{0.55 \pm CH}}{2}.$$

And from the monotonicity we know $\lim_{y\to y_1'^-} g_a(y), \lim_{y\to y_1'^-} g_a'(y)$ exist ($\lim_{y\to y_1'^-} g_a'(y)$ is possibly $-\infty$) and

$$\lim_{y\to y_1'^-} g_a(y) \geq 0.$$

We will see $\lim_{y\to y_1'^-} g_a'(y)$ can not be finite. Should it be finite, if $\lim_{y\to y_1'^-} g_a(y) = 0$, from (13) one could find $y_1 < y_1'$ such that $g_a''(y_1) = 0$ (where $k_a \approx \frac{1}{2}$) which is a contradiction. If $\lim_{y\to y_1'^-} g_a'(y)$ should be finite and $\lim_{y\to y_1'^-} g_a(y) > 0$, as y_1' is finite, from Lemma 3.2 and Lemma 3.1, the solution can be extended beyond y_1 which is also a contradiction. So we have

$$\lim_{y\to y_1^-} g_a'(y) = -\infty.$$

$\square$

Lemma 3.7. *If (H1) or (H2) happens, there exists* $y_3 < +\infty$ *such that* g_a *can be extended to* y_3 *and* $g_a(y_3) > g_a(y_2), g_a'(y_3) > 0, g_a''(y_3) = 0$. *In* $[y_2, y_3]$ *both* g_a *and* g_a' *are monotonically increasing. Moreover, there is* $C > 0$ *which does not depend on* a *such that*

$$\tau(g_a(y_3), g_a'(y_3)) \in [(1-CH)\tau(g_a(y_1), g_a'(y_1)), (1+CH)\tau(g_a(y_1), g_a'(y_1))].$$

Proof. So from (18) we know

$$\frac{d\tau}{dg_a^2} = \frac{1}{2}\rho. \tag{20}$$

From $|\rho| \leq \hat{C}H$, we have

$$|\tau(g_a(y_2), g_a'(y_2)) - \tau(g_a(y_1), g_a'(y_1))| \leq CHg_a(y_1)^2. \tag{21}$$

We know

$$\tau(g_a(y_1), g_a'(y_1)) = (\frac{1}{k_a(y_1)} - 1)g_a(y_1)^2 \in [(1 - CH)g_a(y_1)^2, (1 + CH)g_a(y_1)^2].$$

So

$$\tau(g_a(y_2), g_a'(y_2))) \in [(1 - CH)g_a(y_1)^2, (1 + CH)g_a(y_1)^2]. \tag{22}$$

From $g_a(y_2) \in (0, \frac{1}{2})$ and $g_a'(y_2) = 0$ we can get finer estimate for $g_a(y_2)$, i.e.

$$g_a(y_2) \in ((1 - CH)g_a(y_1)^2, (1 + CH)g_a(y_1)^2).$$

From Lemma 3.1, g_a can be extended beyond y_2. And in a small interval $(y_2, y_2 + \delta)$, $g_a(y) > 0, g_a'(y) > 0, g_a''(y) > 0$. We want to show there is $y_3 > y_2$ such that $g_a''(y_3) = 0$ and in (y_2, y_3), $g_a''(y) > 0$. If it were false, then as long as g_a can be extended and $y > y_2$ we would have $g_a(y) > 0, g_a'(y) > 0, g_a''(y) > 0$. So when $y > y_2$, both g_a and g_a' would be monotonically increasing. Suppose g_a can be extended to $(y_2, \tilde{y}_3)$ but not until $\tilde{y}_3$, where $\tilde{y}_3$ is possibly $+\infty$. However, as $k_a(y_2) = g_a(y_2) \leq (1 + CH)g_a(y_1)^2 \leq (1 + CH)C_4^2 \leq \frac{1}{4}$, $\{(g_a, g_a'); y > y_2\}$ can not intersect with $k_a = \frac{2}{3}$ or one would find a point y_3 such that $g_a''(y_3) = 0$ (where $k_a \approx \frac{1}{2}$). When $y \to \tilde{y}_3^-$, there is a limit for (g_a, g_a') which lies in the region bounded by $g_a = g_a(y_2), g_a' = 0$ and $k_a = \frac{2}{3}$. As in the last lemma, it is easy to prove that $\tilde{y}_3$ has upper bound and from Lemma 3.2 and Lemma 3.1, we can extend the solution beyond $\tilde{y}_3$ which is a contradiction. So there must be a $y_3 > y_2, y_3 < +\infty$ such that $g_a''(y_3) = 0$. We can choose y_3 as the minimum of such values, so in (y_2, y_3), $g_a(y) > 0, g_a'(y) > 0, g_a''(y) > 0$.

From (20) we know

$$|\tau(g_a(y_2), g_a'(y_2)) - \tau(g_a(y_3), g_a'(y_3))| \leq CHg_a^2(y_3).$$

So we have

$$g_a(y_2) \in ((1 - CH)g_a(y_3)^2, (1 + CH)g_a(y_3)^2).$$

Together with (22) we can deduce

$$g_a(y_3) \in [(1 - CH)g_a(y_1), (1 + CH)g_a(y_1)]$$

and

$$\tau(g_a(y_3), g_a'(y_3)) \in [(1 - CH)\tau(g_a(y_1), g_a'(y_1)), (1 + CH)\tau(g_a(y_1), g_a'(y_1))]. \tag{23}$$

□

Lemma 3.8. *If (H1) or (H2) happens,* g_a *can be extended to* $y_4 > y_3$ *and* $g_a(y_4) \in (\frac{1}{2}, 1.2), g_a'(y_4) = 0$*. In* $[y_3, y_4]$*,* g_a *is monotonically increasing and* g_a' *is monotonically decreasing. There is* $C > 0$ *such that*

$$|\tau(g_a(y_4), g_a'(y_4)) - \tau(g_a(y_3), g_a'(y_3))| \leq CH.$$

Proof. From Lemma 3.1, g_a can be extended beyond y_3. As long as g_a can be extended and $y > y_3, g_a > 0, g_a' > 0$, we are going to prove that $g_a'' < 0$. We know in a small interval $(y_3, y_3 + \delta)$ we have $g_a > 0, g_a' > 0$. From (15) we know, when $y > y_3$, as long as the solution could be extended and $g_a > 0, g_a' > 0, k_a < \frac{2}{2+\hat{C}H}$, k_a will monotonically increase. If $k_a > \frac{1}{2-\hat{C}H}$, we have

$$g_a'' = \frac{1}{g}(1 + g_a'^2)(1 - (2+\rho)k_a) < 0.$$

We are going to prove that when $y > y_3, g_a > 0, g_a' > 0, \frac{1}{2+\rho(y_3)} \leq k_a \leq \frac{1}{2-\hat{C}H}$ we still have $g_a'' < 0$. From (23) and $\frac{1}{2+\rho(y_3)} \leq k_a \leq \frac{1}{2-\hat{C}H}$ we know $g_a < C_4$ and $g_a' \geq 1$. Similarly, we can derive (16) and (17) again. So we have $g_a''(y) < 0$. So when $y > y_3, g_a > 0, g_a' > 0$, we have g_a is monotonically increasing and g_a' is monotonically decreasing. So in this interval, g_a can be regarded as a decreasing function of g_a'. We claim that, when $y > y_3$ and $g_a > 0, g_a' > 0$, as long as the solution could be extended,

$$k_a \leq 3. \tag{24}$$

To see this, by contradiction, we assume g_a can be extended to some $\tilde{y}_4 > y_3$ such that when $y \in (y_3, \tilde{y}_4)$ we have $g_a(y) > 0, g_a'(y) > 0$, and $k_a(\tilde{y}_4) > 3$. Then there must be $y_4' \in (y_3, \tilde{y}_4)$ such that $k_a(y_4') = 3$. However if we integral (15) from y_4' to $\tilde{y}_4$ we can get a contradiction. So we proved that $k_a \leq 3$. So we have $g_a \leq 3$. From the monotonicity of g_a and g_a' and Lemma 3.2, obviously the solution can be extended to some $y_4 > y_3$ such that $g_a(y_4) \leq 3, g_a'(y_4) = 0$.

From (20) we know

$$|\tau(g_a(y_4), g_a'(y_4)) - \tau(g_a(y_3), g_a'(y_3))| \leq CH. \tag{25}$$

□

Lemma 3.9. *If (H1) or (H2) happens, there holds*

$$|\tau(g_a(y_4), g_a(y_4)) - (a - a^2)| + |g_a(y_4) - a| \leq CH.$$

Moreover, if we choose $a \in [0.95, 1.05]$ *and* H *very small we have* $g_a(y_4) \in [0.9, 1.1]$.

Proof. From (19)(23)(25) we get the conclusion easily.

□

Lemma 3.10. *If (H1) happens,* g_a *can be extended to* $y_5 > y_4$ *such that* $g_a(y_5) > 0, g_a'(y_5) < 0, g_a''(y_5) = 0$ *and*

$$|\tau(g_a(y_5), g_a'(y_5)) - \tau(g_a(y_4), g_a'(y_4))| \leq CH.$$

If (H2) happens, g_a can be extended to $y_5 > y_4$ such that

$$\lim_{y \to y_5^-} g_a \geq 0,$$
$$\lim_{y \to y_5^-} g'_a = -\infty.$$

In either case, g_a and g'_a are monotonically decreasing in (y_4, y_5).

Proof. This lemma can be proved in the same way as Lemma 3.6.

□

4 Analysis of the singularities

Define a subset $E(H)$ of $[0, 95, 1.05]$ which includes all the a values such that (H1) happens to g_a. And denote $\bar{E}(H) = \{a \in [0.95, 1.05] : a \notin E(H)\}$. Obviously, we can choose H sufficiently small such that $0.95 \in E(H), 1.05 \in \bar{E}(H)$. And from the continuous dependence of the solution of ODE on its initial values, $E(H)$ is an open subset of $[0, 95, 1.05]$. Let

$$a'(H) = \inf \bar{E}(H).$$

We have $a'(H) \in \bar{E}(H)$ and $a'(H) > 0.95$ for small H.

Lemma 4.1.
$$\lim_{a \to a'(H)^-} \max\{|g'_a(y_1)|, |g'_a(y_5)|\} = +\infty.$$

Proof. If it were not true, we could find a sequence a_n which tends to $a'(H)^-$ such that for any n

$$\max\{|g'_{a_n}(y_1)|, |g'_{a_n}(y_5)|\} \leq C(H).$$

So $(g_{a_n}, g'_{a_n}) \in D(A_1(H), A_2(H), A_3(H), A_4(H))$ from Lemma 3.7, 3.8, 3.9, 3.10. So

$$\|g_{a_n}(y)\|_{C^2_y([0,y_5])} \leq C(H). \tag{26}$$

Note that $y_i, i = 1, \cdots, 5$ depend on a and H. Set $y_0 = 0$. We are going to prove that there is $C(H) > 0$ such that

$$C(H)^{-1} \leq y_i - y_{i-1} \leq C(H). \tag{27}$$

We only prove this for $y_1 - y_0$ and $y_2 - y_1$. The rest inequalities follow similarly. It is easy to see that $k_a < 3$ and from the same argument as in the proof of Lemma 3.6 we see when $-\frac{1}{2} < g'_a \leq 0$ on both sides of y_1, $k_a - \frac{1}{2}$ does not change sign and is bounded away from 0. By dividing the orbit into $-\frac{1}{2} < g'_a \leq 0$ and $g'_a \leq -\frac{1}{2}$, we can apply Lemma 3.3 to get (27).

By passing to a subsequence, we may assume that for fixed H, when $a_n \to a'(H)^-$ we have $y_i \to y_i^*$ for $i = 1, \cdots, 5$ and

$$C(H)^{-1} \leq y_i^* - y_{i-1}^* \leq C(H).$$

Differentiate the ODE (6) with respect to y once. From (26) and (16) we know

$$\|g_{a_n}\|_{C^3_y([0,y_5])} \leq C(H).$$

So g_{a_n} converges to some $\tilde{g}(y), y \in [0, y_5^* - \delta]$ in C^2 for any small $\delta > 0$. We know $\tilde{g}(y)$ solves ODE (6) with $a = a'(H)$. From the uniqueness of the ODE, we know $g_{a'(H)}(y) = \tilde{g}(y), y \in [0, y_5^*)$. By uniform C^3 estimate of g_{a_n}, we know $g_{a'(H)}$ can be extended to $y = y_5^*$ and $g''_{a'(H)}(y_5^*) = 0$ and in each interval $[y_{i-1}^*, y_i^*]$ it has the same monotonicity as g_a in $[y_{i-1}, y_i]$. Then from the proof of Lemma 3.5, we know (H1) happens to $g_{a'(H)}$ which is a contradiction. So we proved this lemma. $\square$

So there is $\delta_1(H) > 0$ which depends on H such that when $-\delta_1(H) < a - a'(H) < 0$,

$$\max\{|g'_a(y_1)|, |g'_a(y_5)|\} \geq \frac{1}{\sqrt{H}}.$$

Lemma 4.2. *There is $C > 0$ which does not depend on a, H, such that when $-\delta_1(H) < a - a'(H) < 0$,*

$$\min\{|g'_a(y_1)|, |g'_a(y_5)|\} \geq \frac{C}{\sqrt{H}}.$$

Proof. Let's first assume

$$\frac{1}{\sqrt{H}} \leq \max\{|g'_a(y_1)|, |g'_a(y_5)|\} = |g'_a(y_1)|.$$

From Lemma 3.7 and $g_a(y_1)\sqrt{1 + g'_a(y_1)^2} \in [\frac{1}{2+\hat{C}H}, \frac{1}{2-\hat{C}H}]$ we know there is a uniform $C > 0$ such that

$$\tau(g_a(y_1), g'_a(y_1)), \tau(g_a(y_3), g'_a(y_3)) \in [C^{-1}H, CH]. \tag{28}$$

From Lemma 3.8 and Lemma 3.10, we know

$$0 < \tau(g_a(y_5), g'_a(y_5)) \leq CH.$$

Combining this with $g_a(y_5)\sqrt{1 + g'_a(y_5)^2} \in [\frac{1}{2+\hat{C}H}, \frac{1}{2-\hat{C}H}]$, we know there is $C > 0$ such that

$$|g'_a(y_5)| \geq \frac{C}{\sqrt{H}}.$$

If

$$\max\{|g'_a(y_1)|, |g'_a(y_5)|\} = |g'_a(y_5)|$$

we can prove the lemma similarly. $\square$

Let $h(y) = \sqrt{1-(y-y_4)^2}, y \in [y_4 - 1, y_4 + 1]$. $h(y)$ solves the following ODE

$$\begin{cases} h''(y) - \frac{1}{h(y)}(1 + h'(y)^2) + 2(1 + h'(y)^2)^{\frac{3}{2}} = 0, \\ h(y_4) = 1, \\ h'(y_4) = 0. \end{cases}$$

There is exactly one $y_6 \in (y_3, y_4)$ such that $g_a'(y_6) = 1$. Define two continuous maps

$$\Phi_1 : [y_6, y_4] \to \mathbb{R},$$
$$\Phi_2 : [y_3, y_6] \to \mathbb{R}$$

such that $g_a'(y) = h'(\Phi_1(y))$, $\Phi_1(y_4) = y_4$ and $g_a(y) = h(\Phi_2(y))$, $\Phi_2(y_6) \in (y_4 - 1, y_4)$. We use a graph below to illustrate the definition of Φ_1, Φ_2. In this graph, the horizontal direction represents $g_a'(y)$ or $h'(y)$ and the vertical direction represents $g_a(y)$ or $h(y)$.



We have the following estimates,

Lemma 4.3. *There is $C > 0$ such that when H is very small and $-\delta_1(H) < a - a'(H) < 0$,*

$$|\Phi_1(y) - y| \le CH,$$
$$|\Phi_2(y) - y| \le CH(\frac{H}{g_a^2(y)} + |\log g_a(y)|).$$

Proof.

$$\frac{d\Phi_1(y)}{dy} = \frac{dg_a'(y)}{dy}\frac{d\Phi_1(y)}{dh'(\Phi_1(y))}$$
$$= \frac{\frac{1}{g_a}(1 + g_a'^2) - (2+\rho)(1 + g_a'^2)^{\frac{3}{2}}|_y}{\frac{1}{h}(1 + h'^2) - 2(1 + h'^2)^{\frac{3}{2}}|_{\Phi_1(y)}}.$$

When $-\delta_1(H) < a - a'(H) < 0$, $|\tau(g_a(y), g_a'(y))| \le CH, y \in [0, y_5]$. So we know when $y \in [y_6, y_4]$, $g_a(y) \in [h(\Phi_1(y))(1 - CH), h(\Phi_1(y))(1 + CH)]$. Hence,

$$|\frac{\frac{1}{g_a}(1 + g_a'^2) - (2+\rho)(1 + g_a'^2)^{\frac{3}{2}}|_y}{\frac{1}{h}(1 + h'^2) - 2(1 + h'^2)^{\frac{3}{2}}|_{\Phi_1(y)}} - 1| \le CH.$$

So the first estimate follows. For the second inequality, first we notice

$$|h(\Phi_1(y_6)) - h(\Phi_2(y_6))| = |h(\Phi_1(y_6)) - g_a(y_6)| \leq CH.$$

From $h'(\Phi_1(y_6)) = 1$, we know $\Phi_1(y_6) = y_4 - \frac{\sqrt{2}}{2}$. So $h(\Phi_1(y_6)) = \frac{\sqrt{2}}{2}$ and

$$|\Phi_2(y_6) - \Phi_1(y_6)| \leq CH.$$

When $y \in [y_3, y_6]$,

$$\begin{aligned} y - y_6 &= \int_{g_a(y_6)}^{g_a(y)} \frac{1}{g_a'} dg_a \\ &= \int_{g_a(y_6)}^{g_a(y)} \frac{\tau + g_a^2}{\sqrt{g_a^2 - (\tau + g_a^2)^2}} dg_a \end{aligned}$$

and

$$\begin{aligned} \Phi_2(y) - \Phi_2(y_6) &= \int_{h(\Phi_2(y_6))}^{h(\Phi_2(y))} \frac{1}{h'} dh \\ &= \int_{g_a(y_6)}^{g_a(y)} \frac{g_a}{\sqrt{1 - g_a^2}} dg_a. \end{aligned}$$

So we have

$$\begin{aligned} &\Phi_2(y) - y - (\Phi_2(y_6) - y_6) \\ =& \int_{g_a(y_6)}^{g_a(y)} \left(\frac{g_a}{\sqrt{1 - g_a^2}} - \frac{\tau + g_a^2}{\sqrt{g_a^2 - (\tau + g_a^2)^2}}\right) dg_a \\ =& \int_{g_a(y_6)}^{g_a(y)} \frac{-\tau^2 - 2\tau g_a^2}{\sqrt{1 - g_a^2}\sqrt{g_a^2 - (\tau + g_a^2)^2}(g_a\sqrt{g_a^2 - (\tau + g_a^2)^2} + (\tau + g_a^2)\sqrt{1 - g_a^2})} dg_a. \end{aligned}$$

When $y \in [y_3, y_6]$,

$$\begin{aligned} 1 - g_a^2 &\geq 1 - \left(\frac{\sqrt{2}}{2} + CH\right)^2 \geq \frac{1}{3}, \\ |\tau| &\leq CH. \end{aligned}$$

From (24) we know when $y \in [y_3, y_4]$

$$\frac{\tau + g_a^2}{g_a^2} = \frac{1}{k_a} \geq \frac{1}{3}.$$

So

$$\tau + g_a^2 \geq \frac{g_a^2}{3}. \tag{29}$$

From $g_a \leq \frac{\sqrt{2}}{2} + CH, |\tau| \leq CH$, one can check that, for some $C > 0$,

$$g_a^2 - (\tau + g_a^2)^2 \geq C g_a^2.$$

So we have

$$|\frac{-\tau^2-2\tau g_a^2}{\sqrt{1-g_a^2}\sqrt{g_a^2-(\tau+g_a^2)^2}(g_a\sqrt{g_a^2-(\tau+g_a^2)^2}+(\tau+g_a^2)\sqrt{1-g_a^2})}|\leq C\frac{H^2+Hg_a^2}{g_a^3}.$$

Then

$$\begin{aligned}|\Phi_2(y)-y-(\Phi_2(y_6)-y_6)| &\leq C|\int_{g_a(y_6)}^{g_a(y)}\frac{H^2+Hg_a^2}{g_a^3}dg_a| \\ &= C(|\frac{H^2}{g_a^2(y)}-\frac{H^2}{g_a^2(y_6)}|+H|\log g_a(y)-\log g_a(y_6)|) \\ &\leq CH(\frac{H}{g_a^2(y)}+|\log g_a(y)|).\end{aligned}$$

So we have when $y\in[y_3,y_6]$

$$\begin{aligned}|\Phi_2(y)-y| &\leq |\Phi_2(y_6)-\Phi_1(y_6)|+|\Phi_1(y_6)-y_6|+CH(\frac{H}{g_a^2}+|\log g_a|) \\ &\leq CH(\frac{H}{g_a^2}+|\log g_a|+1).\end{aligned}$$

As $1\leq C|\log\frac{\sqrt{2}}{2}|$, we get the second estimate.

$\square$

Lemma 4.4. *When H is very small and $-\delta_1(H)<a-a'(H)<0$,*

$$|y_4-2|\leq C\sqrt{H},$$
$$|y_5-3|\leq C\sqrt{H}.$$

Proof. From (23) we know

$$h(\Phi_2(y_3))=g_a(y_3)\leq C\sqrt{H}.$$

So there is some $y_6'\in(y_{3,}y_6)$ such that $g_a(y_6')=C\sqrt{H}$ where we use the same constant C as the right hand side of the above inequality. From Lemma 4.3, we know

$$|y_6'-\Phi_2(y_6')|\leq CH(\frac{H}{g_a^2(y_6')}+\log g_a(y_6'))\leq CH|\log H|.$$

We know that $|\Phi_2(y_6')-(y_4-1)|\leq CH$. So we have

$$|y_6'-(y_4-1)|\leq CH|\log H|.$$

By applying Lemma 3.3 we can get

$$|y_6'-y_2|\leq C\sqrt{H}.$$

In the same way we can find $\hat{y}_1 \in (1, y_1)$ such that $g_a(\hat{y}_1) = C\sqrt{H}$ for some $C > 0$. And we can prove that

$$\begin{aligned} |\hat{y}_1 - 1| &\leq CH \log H, \\ |\hat{y}_1 - y_2| &\leq C\sqrt{H}. \end{aligned}$$

So

$$\begin{aligned} |y_4 - 2| &= |y_4 - 1 - y_6' + y_6' - y_2 + y_2 - \hat{y}_1 + \hat{y}_1 - 1| \\ &\leq CH \log H + C\sqrt{H} \leq C\sqrt{H}. \end{aligned}$$

The second estimate follows similarly.

$\square$

Lemma 4.5. *When H is very small and $-\delta_1(H) < a - a'(H) < 0$,*

$$|\int_{y_3}^{y_4} \frac{1}{\sqrt{y^2 + g_a(y)^2}} g_a g_a' dy - \int_{y_4-1}^{y_4} \frac{1}{\sqrt{y^2 + h(y)^2}} hh' dy| \leq CH,$$

$$|\int_{y_3}^{y_4} \frac{g_a - y g_a'}{(y^2 + g_a(y)^2)^{\frac{3}{2}} \sqrt{1 + g_a'^2}} g_a g_a' dy - \int_{y_4-1}^{y_4} \frac{h - yh'}{(y^2 + h(y)^2)^{\frac{3}{2}} \sqrt{1 + h'^2}} hh' dy| \leq CH.$$

Proof. For the first one we have an obvious reason to prove

$$|\int_{y_6'}^{y_4} \frac{1}{\sqrt{y^2 + g_a(y)^2}} g_a g_a' dy - \int_{\Phi_2(y_6')}^{y_4} \frac{1}{\sqrt{y^2 + h(y)^2}} hh' dy| \leq CH$$

instead, where $y_6' \in (y_3, y_6)$ and $g_a(y_6') = C\sqrt{H}$. By using Lemma 4.3, this can be verified as

$$\begin{aligned} &|\int_{y_6'}^{y_6} \frac{1}{\sqrt{y^2 + g_a(y)^2}} g_a g_a' dy - \int_{\Phi_2(y_6')}^{\Phi_2(y_6)} \frac{1}{\sqrt{y^2 + h(y)^2}} hh' dy| \\ \leq &\int_{g_a(y_6')}^{g_a(y_6)} |\frac{g_a}{\sqrt{y^2 + g_a(y)^2}} - \frac{g_a}{\sqrt{\Phi_2(y)^2 + g_a(y)^2}}| dg_a \\ \leq &C \int_{g_a(y_6')}^{g_a(y_6)} g_a H (\frac{H}{g_a^2} + |\log g_a|) dg_a \\ \leq &CH^2 |\log H| + CH \leq CH, \end{aligned}$$

$$|\int_{\Phi_2(y_6)}^{\Phi_1(y_6)} \frac{1}{\sqrt{y^2 + h(y)^2}} hh' dy| \leq CH,$$

$$\begin{aligned} &|\int_{\Phi_1(y_6)}^{y_4} \frac{1}{\sqrt{y^2 + h(y)^2}} hh' dy - \int_{y_6}^{y_4} \frac{1}{\sqrt{y^2 + g_a(y)^2}} g_a g_a' dy| \\ \leq &|\int_{y_6}^{y_4} \frac{1}{\sqrt{\Phi_1(y)^2 + h(\Phi_1(y))^2}} h(\Phi_1(y)) h'(\Phi_1(y)) \Phi_1'(y) - \frac{1}{\sqrt{y^2 + g_a(y)^2}} g_a g_a' dy| \\ \leq &CH. \end{aligned}$$

The proof of the second one is similar.

$\square$

From Lemma 4.2, using the same technique as the above lemma, we can prove

Lemma 4.6. *When H is very small and $-\delta_4(H) < a - a'(H) < 0$, we have*

$$|\int_{y_4}^{y_5} \frac{1}{\sqrt{y^2+g_a(y)^2}} g_a g_a' dy - \int_{y_4}^{y_4+1} \frac{1}{\sqrt{y^2+h(y)^2}} hh' dy| \le CH,$$

$$|\int_{y_4}^{y_5} \frac{g_a - y g_a'}{(y^2+g_a(y)^2)^{\frac{3}{2}}\sqrt{1+g_a'^2}} g_a g_a' dy - \int_{y_4}^{y_4+1} \frac{h - yh'}{(y^2+h(y)^2)^{\frac{3}{2}}\sqrt{1+h'^2}} hh' dy| \le CH.$$

Lemma 4.7. *We can fix $\lambda > 0$ small and $p > 0$ large such that there is $\delta(p,\lambda) > 0$ small such that when $0 < H < \delta(p,\lambda)$ and $-\delta_1(H) < a - a'(H) < 0$, we have $|g_a'(y_5)| \ge |g_a'(y_1)|$. And moreover*

$$\lim_{a\to a'(H)^-} g_a'(y_5) = -\infty, \ \liminf_{a\to a'(H)^-} g_a'(y_1) > -\infty.$$

Proof. From Lemma 4.5 and Lemma 4.6 we know

$$\begin{aligned}
&|\int_{y_3}^{y_5} (\frac{1}{\sqrt{y^2+g_a(y)^2}} + \frac{(g_a - y g_a')}{(y^2+g_a(y)^2)^{\frac{3}{2}}\sqrt{1+g_a'^2}}) g_a g_a' dy \\
&- \int_{y_4-1}^{y_4+1} (\frac{1}{\sqrt{y^2+h(y)^2}} + \frac{h - yh'}{(y^2+h(y)^2)^{\frac{3}{2}}\sqrt{1+h'^2}}) hh' dy| \\
\le& CH.
\end{aligned}$$

The following facts are the key to our proof. When $y_4 > 1$,

$$\int_{y_4-1}^{y_4+1} (\frac{1}{\sqrt{y^2+h(y)^2}} + \frac{h - yh'}{(y^2+h(y)^2)^{\frac{3}{2}}\sqrt{1+h'^2}}) hh' dy = 0,$$

$$\int_{y_4-1}^{y_4+1} \frac{1}{\sqrt{y^2+h(y)^2}} hh' dy = \frac{2}{3y_4^2},$$

$$\int_{y_4-1}^{y_4+1} \frac{h - yh'}{(y^2+h(y)^2)^{\frac{3}{2}}\sqrt{1+h'^2}} hh' dy = -\frac{2}{3y_4^2}.$$

The proof of these facts takes direct calculations which we omit here.
From Lemma 4.4, we know $|y_4 - 2| \le C\sqrt{H}$. So

$$\int_{y_4-1}^{y_4+1} \frac{h - yh'}{(y^2+h(y)^2)^{\frac{3}{2}}\sqrt{1+h'^2}} hh' dy = -\frac{1}{6} \pm O(\sqrt{H}).$$

So we have

$$|\int_{y_3}^{y_5} (\frac{H}{\sqrt{y^2+g_a(y)^2}} + \frac{H(g_a - y g_a')}{(y^2+g_a(y)^2)^{\frac{3}{2}}\sqrt{1+g_a'^2}}) g_a g_a' dy| \le CH^2.$$

and when $p > 3$

$$\int_{y_3}^{y_5} \frac{(p-3)H^2(g_a - yg'_a)}{(y^2+g_a(y)^2)^{\frac{3}{2}}\sqrt{1+g'^2_a}} g_a g'_a dy \le -(p-3)H^2(\frac{1}{6}+CH)$$
$$\le -\frac{(p-3)}{7}H^2,$$

$$|\int_{y_3}^{y_5} \frac{(\phi-p)H^2(g_a - yg'_a)}{(y^2+g_a(y)^2)^{\frac{3}{2}}\sqrt{1+g'^2_a}} g_a g'_a dy| \le C|p-1|H^2\lambda^2,$$

From (8)

$$|\int_{y_3}^{y_5}(-\frac{H^2\phi'(y+g_a g'_a)}{8d^2y^2\sqrt{1+g'^2_a}} + C(p,\lambda)O(H^3))g_a g'_a dy| \le C|p-1|\lambda H^2 + C(p,\lambda)\lambda^2 H^3.$$

Also note that

$$|\int_{y_1}^{y_3} g_a g'_a \rho dy| = |\int_{g_a(y_1)}^{g_a(y_2)} + \int_{g_a(y_2)}^{g_a(y_3)} \frac{1}{2}\rho dg_a^2|$$
$$\le CH^2.$$

So we can fix $\lambda > 0$ small and $p > 0$ large, and there is $\delta(\lambda, p) > 0$ small that when $0 < H < \delta(\lambda, p)$

$$-CpH^2 \le \tau(g_a(y_5), g'_a(y_5)) - \tau(g_a(y_1), g'_a(y_1)) = \int_{y_1}^{y_5} \rho g_a g'_a dy \le -H^2,$$
$$-CpH^2 \le \tau(g_a(y_5), g'_a(y_5)) - \tau(g_a(y_3), g'_a(y_3)) = \int_{y_3}^{y_5} \rho g_a g'_a dy \le -H^2. \quad (30)$$

From $\tau(g_a(y_i), g'_a(y_i)) > 0, i = 1, 3, 5$, we know

$$\tau(g_a(y_i), g'_a(y_i)) \ge H^2, i = 1, 3.$$

Note that $k_a(y_i) = \frac{1}{2\pm O(H)}, i = 1, 3, 5$. So

$$g'_a(y_i) = \frac{-\sqrt{\tau - 4\tau^2}}{2\tau} = -\frac{1}{2\sqrt{\tau}}(1+O(\tau)), i = 1, 5,$$
$$g'_a(y_3) = \frac{1}{2\sqrt{\tau}}(1+O(\tau)).$$

So we have

$$0 > g'_a(y_1) \ge -\frac{C}{H},$$
$$0 < g'_a(y_3) \le \frac{C}{H},$$

and

$$\lim_{a \to a'(H)^-} g'_a(y_5) = -\infty.$$

□

Now we can analyze the singular limit of g_a as $a \to a'(H)^-$.

Lemma 4.8. *Choose* $p, \lambda, \delta(p, \lambda), \delta_1(H)$ *such that the conclusion of Lemma 4.7 holds. When* $a \to a'(H)^-$*, by passing to a sequence, we have* $y_i \to y_i^*, i = 1, \cdots, 5$ *and* g_a *converges to* $g_{a'(H)}$ *in* C^2 *sense on every* $[0, \tilde{y}] \subset [0, y_5^*)$*. (H2) happens to* $g_{a'(H)}(y)$ *and*

$$\lim_{y \to y_5^{*-}} g_{a'(H)}(y) = 0,$$

$$\lim_{y \to y_5^{*-}} g'_{a'(H)}(y) = -\infty.$$

Proof. From

$$\lim_{a \to a'(H)^-} g'_a(y_5) = -\infty$$

and g'_a is monotonically decreasing on $[y_4, y_5]$ we know for any $j \in \mathbb{N}^+$ when a is sufficiently close to $a'(H)$, there is exactly one $y_5^j \in (y_4, y_5)$ such that $g'_a(y_5^j) = -j$. From

$$-\frac{C}{H} \leq g'_a(y_1) < 0 < g'_a(y_3) \leq \frac{C}{H}$$

we know that g'_a is bounded independent of a on $[0, y_5^j]$. From the analysis of the last section, it is not hard to prove that g_a has both positive upper bound and positive lower bound on $[0, y_5^j]$. So $(g_a, g'_a) \in D(A_1, A_2, A_3, A_4)$, where $A_i = A_i(H, j), i = 1, \cdots, 4$. So on $[0, y_5^j]$, g_a has uniform C^3 bounds. By using the same argument as in Lemma 4.1, we can prove that there is $C(p, \lambda, H) > 0$ such that

$$C(p, \lambda, H)^{-1} \leq y_i - y_{i-1} \leq C(p, \lambda, H), i = 1, \cdots, 4.$$

And from Lemma 3.3, we can prove that $C^{-1} < y_5^j - y_4 < C$. So by passing to a sequence, we can assume $y_i \to y_i^*, i = 1, \cdots, 4, y_5^j \to y_5^{j*}$. And g_a converges to $g_{a'(H)}$ in $C^2([0, y_5^{j*}])$ sense, where $g_{a'(H)}$ solves (6) with initial value $a'(H)$. By a diagonal method, we can choose $a_i \to a'(H)^-$ such that for any $j \in \mathbb{N}^+$

$$\lim_{i \to +\infty} y_5^j = y_5^{j*}.$$

It is obvious that y_5^{j*} is monotonically increasing in j. Note that we have

$$y_5^{j+1} - y_5^j = \int_{-(j+1)}^{-j} \frac{g_a dg'_a}{(1 + g_a'^2)((2 + \rho)k_a - 1)}.$$

For any $j > 0$, when $-j - 1 < g'_a < -j$, we can choose a close to $a'(H)$ such that k_a is bounded away from (and bigger than) $\frac{1}{2}$. So there is $C_j > 0$ such that

$$y_5^{j+1} - y_5^j \geq C_j.$$

So we know $|y_5^{j+1*} - y_5^{j*}| \geq C_j$. From the fact that the second derivative of g_a has uniform bound in $[0, y_5^j]$, we know

$$g'_{a'(H)}(y_5^{j*}) = \lim_{a \to a'(H)^-} g'_a(y_5^{j*}) = \lim_{a \to a'(H)^-} g'_a(y_5^j) = -j.$$

So we know
$$\lim_{j\to+\infty} g'_{a'(H)}(y_5^{j*}) = -\infty.$$

Note that
$$|y_5^j - y_5| \le \frac{C}{j}.$$

If we take a subsequence further, we have $y_5 \to y_5^*$. So we have $|y_5^{j*} - y_5^*| \le \frac{C}{j}$. So we have
$$\lim_{j\to+\infty} y_5^{j*} = y_5^*.$$

From the monotonicity of g'_a we know
$$\lim_{y\to y_5^*} g'_{a'(H)}(y) = -\infty.$$

For $k_a = g_a\sqrt{1+g_a'^2}$, by analyzing (15), we know when $y \in [y_4, y_5]$,
$$\frac{1}{2+CH} \le k_a \le \frac{1}{1-CH}. \tag{31}$$

So we have
$$0 < g_a(y_5^j) \le \frac{C}{j}$$

hence $g_{a'(H)}(y_5^{j*}) \le \frac{C}{j}$ and
$$\lim_{y\to y_5^*} g_{a'(H)}(y) = 0.$$

At last, noticing that any $[0,\tilde{y}] \subset [0, y_5^*)$ is contained in some $[0, y_5^j)$ for any a close to $a'(H)$. So we can prove the uniform convergence on $[0,\tilde{y}]$. Obviously, (H2) happens to $g_{a'(H)}$.

$\square$

Here we draw a graph of $g_{a'(H)}(y), y \in [0, y_5^*)$.



Singularity develops at y_5

$g_{a'(H)}(y)$ is monotonically decreasing in $[y_4^*, y_5^*)$. Now we use g to represent $g_{a'(H)}$ for short. Suppose $y = y(g), g \in (0, g_{a'(H)}(y_4^*)]$ is the inverse function of

$g_{a'(H)}(y), y \in [y_4^*, y_5^*)$. We make even extension to $y(g)$.

$$y^{even}(g) = \begin{cases} y(g), & g \in (0, g_{a'(H)}(y_4^*)], \\ y_5^*, & g = 0, \\ y(-g), & g \in [-g_{a'(H)}(y_4^*), 0). \end{cases}$$

Then we have

Lemma 4.9. *$y^{even}(g)$ has continuous second derivative in $(-\delta, \delta)$ for small δ.*

Proof. $y^{even}(g)$ is continuous at 0. When $g > 0$,

$$\frac{dy^{even}}{dg} = \frac{1}{g'}.$$

From (31), we know when g is close to 0, $g' \sim \frac{1}{g}$. So we have

$$\lim_{g \to 0^+} \frac{dy^{even}}{dg} = 0.$$

So $y^{even}(g)$ has 0 derivative at $g = 0$. For the second derivative, we note that

$$\begin{aligned} \frac{d^2 y^{even}}{dg^2} &= \frac{d}{dg}(\frac{1}{g'}) = \frac{-g''}{g'^3} \\ &= -\frac{1+g'^2}{g'^2}\frac{1}{g}\frac{1}{g'} + (2+\rho)\frac{(1+g'^2)^{\frac{3}{2}}}{g'^3}. \end{aligned} \tag{32}$$

Regard $\frac{1+g'^2}{g'^2}$ as $G_1(g)$ and $\frac{(1+g'^2)^{\frac{3}{2}}}{g'^3}$ as $G_2(g)$. We get a linear ODE of first order,

$$\begin{cases} \frac{d}{dg}(\frac{1}{g'}) + \frac{G_1(g)}{g}\frac{1}{g'} & = (2+\rho)G_2(g), \\ \frac{1}{g'}(0) = 0, \end{cases}$$

with

$$\begin{aligned} \lim_{g \to 0^+} G_1(g) &= 1, \\ G_1(g) &= 1 + O(g^2), \\ \lim_{g \to 0^+} G_2(g) &= 1. \end{aligned}$$

The solution is

$$\frac{1}{g'} = \frac{\int_0^g (2+\rho)G_2(s)s\exp(O(s^2))ds}{g\exp(O(g^2))}.$$

So

$$\begin{aligned} \lim_{g \to 0^+} \frac{1}{gg'} &= \lim_{g \to 0^+} \frac{\int_0^g (2+\rho)G_2(s)s\exp(O(s^2))ds}{g^2\exp(O(g^2))} \\ &= \lim_{g \to 0^+} \frac{(2+\rho)G_2(g)g\exp(O(g^2))}{2g\exp(O(g^2)) + g^2\exp(O(g^2))O(g)} \\ &= 1 + \frac{1}{2}\lim_{g \to 0^+} \rho. \end{aligned}$$

From (7) we know as $y \to y_5^*, g_{a'(H)} \to 0^+, g'_{a'(H)} \to -\infty$, we have $d \to y_5^*$, $\frac{g_a - y g'_a}{\sqrt{1+g_a'^2}} \to y_5^*$. So the limit

$$\lim_{g \to 0^+} \rho$$

exists. So from (32), $\frac{d^2 y^{even}}{dg^2}(0)$ exists.

□

5 Proof of the main theorems

Now we prove Theorem 1.1. For any H sufficiently small, we have got $g_{a'(H)}(y)$, where we have assume that $g_{a'(H)}(y_5^*) = 0$. Consider the surfaces of revolution $\Sigma^+(H)$ defined by

$$r = \frac{2}{H} g_{a'(H)}(\frac{H}{2}x), x \in [0, \frac{2y_5^*}{H}].$$

From our construction, when $x \in [0, \frac{2y_5^*}{H})$ the surface has constant mean curvature H. When $x = \frac{2y_5^*}{H}$, from Lemma 4.9, the surface actually has well defined and continuous mean curvature at this single point. So the mean curvature at this point must be H. Let $\Sigma^-(H)$ be defined by

$$r = \frac{2}{H} g_{a'(H)}(-\frac{H}{2}x), x \in [-\frac{2y_5^*}{H}, 0).$$

And let $\Sigma(H) = \Sigma^+(H) \cup \Sigma^-(H)$. Then $\Sigma(H)$ has constant mean curvature H globally. It is obvious that $\Sigma(H)$ has sphere topology and it is embedded.

From Lemma 3.4, we know $C^{-1}H^{-1} \le l_0 \le CH^{-1}$. So it follows that for any compact set K, as long as H is sufficiently small, $\Sigma(H)$ separates K from infinity.

Now we calculate $|\Sigma(H)|$. Let $|\Sigma(H)|_e$ denote the area in Euclidean metric. We know that

$$|\frac{|\Sigma(H)|_e}{|\Sigma(H)|} - 1| \le C l_0^{-1} \le CH.$$

So we need only to prove that $|H^2|\Sigma(H)|_e - 48\pi| \le C(p)H$. If we denote $g_{a'(H)}$ as g for short, we have

$$H^2|\Sigma(H)|_e = 16\pi \int_0^{y_5^*} g\sqrt{1+g'^2}dy.$$

First we consider the integral on $[y_2^*, y_4^*]$. From (30) we know,

$$H^2 \le \tau(g(y_3^*), g'(y_3^*)) \le CpH^2$$

Then for some $C(p) > 0$, $CH \le g(y_3^*) \le C(p)H, C(p)H^{-1} \le g'(y_3^*) \le CH^{-1}$.

There is $y_6^* \in [y_3^*, y_4^*]$ such that $g'(y_6^*) = 1$. We have

$$
\begin{aligned}
|\int_{y_6^*}^{y_4^*} g\sqrt{1+g'^2}dy - \int_{\Phi_1(y_6^*)}^{y_4^*} h(s)\sqrt{1+h'^2(s)}ds| &\le C(p)H,\\
|\int_{\Phi_2(y_6^*)}^{\Phi_1(y_6^*)} h(s)\sqrt{1+h'^2(s)}ds| &\le C(p)H,\\
|\int_{g(y_3^*)}^{g(y_6^*)} g\frac{\sqrt{1+g'^2}}{g'}dg - \int_{g(y_3^*)}^{g(y_6^*)} h\frac{\sqrt{1+h'^2}}{h'}dh| &\le C(p)H,\\
|\int_{y_2^*}^{y_3^*} g\sqrt{1+g'^2}dy|, |\int_{y_4^*-1}^{\Phi_2(y_3^*)} h\sqrt{1+h'^2}dy| &\le C(p)H,
\end{aligned}
$$

where the third one holds because

$$
|g(\frac{\sqrt{1+g'^2}}{g'} - \frac{\sqrt{1+h'^2}}{h'})| \le \frac{C(p)H^2}{g} + C(p)gH
$$

and the fourth one holds because $y_3^* - y_2^* \le C(p)H$.

By direct calculation we know

$$
\int_{y_4^*-1}^{y_4^*} h\sqrt{1+h'^2}dy = 1.
$$

So gathering all the inequalities we have

$$
|\int_{y_2^*}^{y_4^*} g\sqrt{1+g'^2}dy - 1| \le C(p)H.
$$

In a similar way we can prove that

$$
|\int_0^{y_2^*} g\sqrt{1+g'^2}dy - 1| \le C(p)H,
$$

$$
|\int_{y_4^*}^{y_5^*} g\sqrt{1+g'^2} - 1| \le CH.
$$

So we know

$$
|\int_0^{y_5^*} g\sqrt{1+g'^2} - 3| \le C(p)H.
$$

So

$$
|H^2|\Sigma(H)|_e - 48\pi| \le C(p)H.
$$

However, if we revise Qing and Tian's proof of the uniqueness CMC spheres in [12], we know for a stable CMC sphere that separates the compact part from infinity and with l_0 large, $H^2|\Sigma|$ should be close to 16π. So the CMC spheres we constructed are unstable.

Now we prove Theorem 1.2. Let's revise Lemma 4.7. Similarly we can choose $p < 0$ small and $\delta(p) > 0$ such that when $0 < H < \delta(p)$ and $-\delta_1(H) < a - a'(H) < 0$, we have $|g'_a(y_5)| < |g'_a(y_1)|$. So we can choose a proper $p = p(H)$ such that $|g'_a(y_5)| = |g'_a(y_1)|$. In this case, we have

$$\lim_{a \to a'(H)^-} g'_a(y_5) = \lim_{a \to a'(H)^-} g'_a(y_1) = -\infty.$$

Now we can analyze the behavior of g_a on y_1 and y_3 by using the same method as used in Lemma 4.8. We will get a singular limit of three spheres (each one is embedded). They share the same axis and the central one meets its two neighbors at two poles where $x_2 = x_3 = 0$. By using the same method as used in Lemma 4.9, we know all the three spheres are smooth and have constant mean curvature H.

The problem is, for different H, we may get different $p(H)$ (which is bounded independent of H). Note that for a particular $H > 0$, when we do all the constructions above, only the metric in a domain

$$D(H) = \{(x_1, x_2, x_3) | C^{-1}H^{-1} \le \sqrt{x_1^2 + x_2^2 + x_3^2} \le CH^{-1}\}$$

really matters. If we have constructed $\Sigma_n^i, i = 1, 2, 3$ for some $H_n > 0$, we can choose a much larger scale to construct $\Sigma_{n+1}^i, i = 1, 2, 3$. We may assume that $D(H_i), D(H_j)$ are disjoint for $i \neq j$. We choose p as a smooth function of $l = \sqrt{x_1^2 + x_2^2 + x_3^2}$ such that $p = p_i$ in each $D(H_i)$. We can assume that $D(H_i)$ is far away from $D(H_{i+1})$ and $p'(l)$ is small enough such that the $\phi_{\lambda,p}$ term does not influence the mass. In this way, we get a smooth asymptotically Schwarzschild metric with mass 1. Then we can prove Theorem 1.2.

References

[1] S. Brendle. Constant mean curvature surfaces in warped product manifolds. *Publ. Math-Paris.*, 117(1):247–269, 2013.

[2] S. Brendle and M. Eichmair. Large outlying stable constant mean curvature spheres in initial data sets. *Invent. Math.*, 197(3):663–682, 2014.

[3] A. Carlotto, O. Chodosh, and M. Eichmair. Effective versions of the positive mass theorem. *arXiv:1503.05910v1*, 2015.

[4] L.H. Huang. Foliations by stable spheres with constant mean curvature for isolated systems with general asymptotics. *Comm. Math. Phys.*, 300(2):331–373, 2010.

[5] G. Huisken and S.T. Yau. Definition of center of mass for isolated physical systems and unique foliations by stable spheres with constant mean curvature. *Invent. Math.*, 124:281–311, 1996.

[6] S. Ma. Uniqueness of the foliation of constant mean curvature spheres in asymptotically flat 3-manifolds. *Pac. J. Math.*, 252(1):145–179, 9 2011.

[7] S. Ma. On the radius pinching estimate and uniqueness of the cmc foliation in asymptotically flat 3-manifolds. *Adv. Math.*, 288:942–984, 2016.

[8] S. Ma and F. Pacard. Constant mean curvature surfaces of delaunay type along a closed geodesic. *arXiv:1412.4240*, 2014.

[9] C. Nerz. Foliations by stable spheres with constant mean curvature for isolated systems without asymptotic symmetry. *arxiv:1408.0752v2*.

[10] A. Neves and G. Tian. Existence and uniqueness of constant mean curvature foliation of asymptotically hyperbolic 3-manifolds. *Geom. Funct. Anal.*, 19(3):910–942, 12 2009.

[11] A. Neves and G. Tian. Existence and uniqueness of constant mean curvature foliation of asymptotically hyperbolic 3-manifolds ii. *J. Reine. Angew. Math.*, 641:69–93, 2010.

[12] J. Qing and G. Tian. On the uniqueness of the foliation of spheres of constant mean curvature in asymptotically flat 3-manifolds. *J. Am. Math. Soc.*, (4):1091–1110, 2007.